FUNGICIDE

ADAMA

SORADUO™

Advanced disease protection powered by Asorbital® Formulation Technology. SORADUO™ provides proven Fusarium protection in wheat and barley.

POWERED BY Asorbital® FORMULATION TECHNOLOGY

GROUP 3

Active Ingredients:

Prothioconazole 250 g/L = EC
and Tebuconazole 430 g/L = SC

Application Rates:

Rate: 121 – 162 ml/ac SORADUO™ A
+ 70 – 94 ml/ac SORADUO™ B

Packaging and Acres Treated:

Co-pack: 1 × 9.71 L SORADUO™ A
+ 1 × 5.65 L SORADUO™ B

Acres Treated: 60 – 80 ac/case

Water Volume:

Ground: 40 – 80 L/ac
(10 – 20 US gal/ac)

Aerial: 20 L/ac (5 US gal/ac)

Rainfastness:

Avoid applying when rain is forecast.

REGISTERED CROPS:

- Barley
- Wheat (durum, spring, winter)

Crop	Disease	Rate	Timing
Barley	For suppression of Fusarium Head Blight (Fusarium spp.)	162 ml/ac SORADUO™ A + 94 ml/ac SORADUO™ B	70 – 100% head emergence to 3 days after full head emergence
Wheat (durum, spring, winter)		121-162 ml/ac SORADUO™ A + 70-94 ml/ac SORADUO™ B	75% head emergence – 50% main stem flower

Maximum 1 application per year.

HOW IT WORKS:

SORADUO™ is a combination of a triazole fungicide that features Asorbital® Formulation Technology and broad-spectrum system activity plus long-lasting foliar protection.

KEY BENEFITS:

- Proven Fusarium protection in Wheat and Barley
- Absorbital® Formulation Technology helps to deliver uniform coverage for consistent protection
- Simple choice without the complication of grower programming
- Application flexibility
- Performance over a wide range of conditions

FUNGICIDE

SORADUO™

REGISTERED AND SUPPORTED TANK MIXES:

Optional: Non-ionic surfactant (NIS) at 0.125% v/v.

MIXING INSTRUCTIONS:

1. ½ fill the tank with clean water.
2. Add required amount of SORADUO™ B.
3. Add required amount of SORADUO™ A.
4. Add optional non-ionic surfactant (NIS).
5. Fill the tank and agitate again before use.

CROP ROTATIONS:

No restrictions.

PRE-HARVEST INTERVAL:

36 days

GRAZING RESTRICTIONS:

6 days

STORAGE:

Do not freeze.

Please note that there has been a re-evaluation of Tebuconazole by the PMRA; the total seasonal rate is limited to a maximum of 136 g a.i./ha/year, except for those crops where the yearly total rate is currently below 136 g a.i./ha.

TIMING AND RATE OF DATA BELOW:

- Sprayed at heading
- SORADUO™ = Prothioconazole 162 ml/ac + Tebuconazole 94 ml/ac
- Prosaro® = 324 ml/ac

FUSARIUM CONTROL IN CEREALS
Summary of 31 trials from 2019-2021
(wheat = 13, barley = 10, durum = 7, winter wheat = 1)



FUSARIUM CONTROL IN CEREALS – YIELDS
Summary of 35 trials from 2019-2021
(wheat = 14, barley = 10, durum = 7, winter wheat = 4)



*Untreated Check

CLICK HERE FOR FULL PRODUCT DETAILS.

ADAMA

Always read and follow label directions.
Toll-free: 1.855.264.6262 | Website: ADAMA.COM

ASORBITAL® is a registered trademark, and SORADUO™ is a trademark of an ADAMA Group Company. All other products are trademarks of their respective companies. © 2026 ADAMA Agricultural Solutions Canada Ltd.

QUICK TIPS:

Fusarium Head Blight outbreaks in wheat and barley occur when the weather is warm and wet at head emergence and flowering. Timing of application is critical when providing protection against Fusarium Head Blight.

2026-01